



# HIV-1 genetic diversity and drug resistance among treatment naïve patients from Southern Brazil: An association of HIV-1 subtypes with exposure categories

Tiago Gräfa, Caroline P.B. Passaesb, Luis G.E. Ferreira<sup>c</sup>, Edmundo C. Grisard<sup>a</sup>, Mariza G. Morgado<sup>b</sup>, Gonzalo Bello<sup>b</sup>, Aguinaldo R. Pinto<sup>a,\*</sup>

<sup>a</sup> Departamento de Microbiologia, Imunologia e Parasitologia, Centro de Ciências Biológicas, Universidade Federal de Santa Catarina, 88040-900 Florianópolis, SC, Brazil

<sup>b</sup> Laboratório de AIDS e Imunologia Molecular, Instituto Oswaldo Cruz, Av. Brasil, 4365 Manguinhos, 21045-900 Rio de Janeiro, RJ, Brazil

<sup>c</sup> Hospital Regional Homero de Miranda Gomes, Rua Adolfo Donato da Silva, Praia Comprida, 88.103-901 São José, SC, Brazil

## ARTICLE INFO

### Article history:

Received 25 January 2011

Received in revised form 13 April 2011

Accepted 25 April 2011

### Keywords:

HIV-1

Molecular epidemiology

Drug resistance mutations

Southern Brazil

## ABSTRACT

**Background:** The AIDS epidemic in Southern Brazil has unique features, showing co-circulation of HIV-1 subtypes C, B and recombinant forms. Florianópolis has the second highest AIDS incidence among Brazilian capitals, but limited information is available about HIV molecular epidemiology and prevalence of primary drug resistance.

**Objectives:** To investigate the molecular epidemiology of HIV-1 in Florianópolis and to describe the prevalence of primary HIV-1 drug resistance mutations (DMRs).

**Study design:** Epidemiological and clinical data from 82 untreated patients from Florianópolis (2008–2009) were analyzed. The HIV-1 subtype at envelope, protease, reverse transcriptase and integrase regions were determined by phylogenetic and bootscanning analyses and the drug resistance profile were analyzed at the Stanford HIV Drug Resistance Database.

**Results:** The most frequent HIV-1 genetic form was subtype C (65.8%) followed by mosaics BC (18.3%), subtype B (13.4%), subtype F1 (1.2%) and BCF1 recombinant (1.2%). HIV-1 subtype C and BC recombinants were much more frequent in the heterosexual exposure category, whereas subtype B was more common in the MSM exposure category. DRMs were seen in 11% of the sequences, 2.4% of them were related to PI, 5% to NRTI, 3.6% to NNRTI and 1.2% was related to INTI.

**Conclusions:** The present study confirms the high prevalence of subtype C and BC recombinants in Santa Catarina State and revealed a significant difference in the subtype distribution among distinct virus exposure categories. This study also shows a relative high prevalence of protease/reverse transcriptase primary drug resistance mutations and corroborates the usefulness of the integrase inhibitors in southern Brazil.

© 2011 Elsevier B.V. Open access under the [Elsevier OA license](http://creativecommons.org/licenses/by-nc-sa/3.0/).

## 1. Background

The HIV/AIDS pandemic has a major impact in Brazil, being the most affected Country in Latin America with over 600,000 people infected.<sup>1</sup> The Brazilian HIV-1 epidemic can be divided in two distinct scenarios. While subtype B prevails in most of the Country, with a secondary occurrence of subtype F1 and BF1

recombinants,<sup>2–9</sup> Southern Brazil presents a distinct pattern with high frequency of subtypes C, B and BC recombinants, following a lower proportion of subtype F1 and BF1 recombinants,<sup>10–17</sup> comprising 19% of the infected individuals in the Country.


The universal access of Brazilian HIV-positive patients to free-of-cost licensed antiretroviral drugs has substantially reduced the AIDS-related mortality.<sup>1</sup> The effectiveness of antiretroviral therapy, however, may be limited by the increasing of resistance to antiretroviral drugs and by transmission of such drug-resistant viruses to drug-naïve patients. National surveys suggest that primary HIV resistance in Brazil is generally low, ranging from 6% to 8%.<sup>2,18,19</sup> However, very high prevalence of HIV primary drug resistance (18–37%) were observed in some localities, such as the City of Santos (Southeastern region)<sup>20,21</sup> and the State of Bahia (North-eastern region),<sup>22</sup> indicating important differences across country regions.

Most of the molecular epidemiology studies on HIV in the Southern region were carried out in the State of Rio Grande do Sul, but

**Abbreviations:** AIDS, acquired immune deficiency syndrome; ARV, antiretroviral; CPR, calibrated population resistance tool; CRF, circulating recombinant form; DRMs, drug resistance mutations; INT, integrase; INTI, integrase inhibitors; HIV, human immunodeficiency virus; MSM, men who have sex with men; NJ, neighbor-joining; NNRTI, non-nucleoside reverse transcriptase inhibitors; NRTI, nucleoside reverse transcriptase inhibitors; PCR, polymerase chain reaction; PI, protease inhibitors; PR, protease; RT, reverse transcriptase; URF, unique recombinant form.

\* Corresponding author. Tel.: +55 48 3721 5206; fax: +55 48 3721 9258.

E-mail address: [pintoar@ccb.ufsc.br](mailto:pintoar@ccb.ufsc.br) (A.R. Pinto).



**Fig. 1.** Geographical location of Florianópolis and frequency of HIV-1 subtypes at the different genomic regions analyzed. (a) Map of Brazil highlighting the state of Santa Catarina (SC) and the metropolitan region of Florianópolis. (b) HIV-1 subtypes were determined by phylogenetic and bootscanning analyses, as described in Section 3.

limited data concerning the prevalence of HIV-1 subtypes and HIV primary drug resistance in the other southern states (Santa Catarina and Paraná) is available. Noteworthy, Santa Catarina State ranks third on AIDS incidence rate in Brazil (29.6 new cases per 100,000 inhabitants in 2008), while its capital, Florianópolis city, occupies the second position in the ranking of capitals (53.7 new cases per 100,000 inhabitants in 2008).<sup>1</sup> Moreover, Florianópolis city is a very important tourist pole in Southern Brazil, receiving people from other South American countries, mainly Argentina and Uruguay, which could have implications on the distribution of HIV subtypes and recombinant forms.

## 2. Objectives

This study describes the HIV-1 genetic diversity among different exposure categories and the prevalence of HIV primary drug resistance mutations based on sequence analyses of protease, reverse transcriptase, integrase and envelope genes among naïve patients from Florianópolis metropolitan area, Santa Catarina, Brazil.


## 3. Study design

### 3.1. Patients


Blood samples were collected from 82 HIV-infected patients followed up at Hospital Homero de Miranda Gomes located in metropolitan region of Florianópolis, Santa Catarina, Brazil. The individuals were selected from May-2008 to February-2009 and inclusion criteria included age over 18-year old, no sign or symptom of AIDS and no previous antiretroviral therapy. The Universidade Federal de Santa Catarina Ethics Committee approved the study and informed consent was obtained from all volunteers.

### 3.2. Extraction, amplification and sequencing of HIV-1 DNA

PBMC was isolated from 4 ml of blood by the Histopaque (Sigma-Aldrich, St. Louis, MO, USA) density gradient method and genomic DNA was extracted using a QIAmp Blood Kit (Qiagen Inc., Chatsworth, CA, USA). HIV-1 protease (PR), reverse transcriptase (RT), integrase (INT) and envelope (*env*) genes were



**Fig. 2.** Phylogenetic analysis of CRF31\_BC-like sequences. (a) Neighbor-joining phylogenetic tree of PR/RT region (2316–3316 nt relative to HXB2), including only the CRF31\_BC-like sequences from Santa Catarina (black diamonds) and subtype reference sequences. Reliability was tested with 1000 replicates bootstrapped, values considering significant (above 70%) are shown. (b) Bootscanning patterns corresponding to a CRF31\_BC reference sequence and the three sequences from Santa Catarina classified as CRF31\_BC-like. Reference sequences were extracted from HIV Los Alamos Database.



**Fig. 3.** Schematic drawing showing breakpoint pattern of the mosaics viruses found in the study. Breakpoint positions were obtained using Simplot 3.5.1 and numbered according to HXB2 reference. The *pol* (PR/RT, INT) and *env* (*gp120*-C2V3) genes are colored according to the subtype: subtype B is red, subtype C is blue and subtype F1 is represented in green. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of the article.)

**Table 1**

General characteristics of HIV-1 infected individuals according to the HIV-1 subtypes.

	Total (n=82)	Subtype C (n=54)	Subtype B (n=11)	BC mosaic (n=15)
Mean age (years)	37.8 ± 10	39 ± 9.2	31.3 ± 9.3	37 ± 11
Sex (%)				
Male	40 (48.8%)	23 (43%)	9 (82%)	7 (46.7%)
Female	42 (51.2%)	31 (57%)	2 (18%)	8 (53.3%)
Mean diagnostic year	2004 ± 3.8	2004 ± 4.3	2005 ± 2.1	2005 ± 2.7
Exposure categories				
Heterosexual	58 (70.6%)	41 (76%)	3 (27.3%)	12 (80%)
MSM	18 (22%)	9 (16.5%)	8 (72.7%)	1 (6.6%)
Injection drug user	4 (5%)	3 (5.5%)	–	1 (6.6%)
Unknown	2 (2.4%)	1 (2%)	–	1 (6.6%)
Mean CD4 T-cell count	513 ± 312	492 ± 269	563 ± 429	522 ± 366
Mean viral load (log <sub>10</sub> copies/ml)	4.1 ± 0.8	4.1 ± 0.8	4.4 ± 0.9	3.9 ± 0.7

Values are expressed ± standard deviation of the mean.

amplified by a nested-PCR as described elsewhere.<sup>23–25</sup> Purified products were sequenced using MegaBace® 1000 DNA Analysis System (GE Healthcare). Sequences were visually inspected and assembled by using SeqMan software, LaserGene package (DNAS-TAR, Madison, WI, USA). Sequences were submitted to GeneBank under accession numbers JF773818–JF774062.

### 3.3. HIV-1 subtype classification

Nucleotide sequences were aligned by using Clustal X program.<sup>26</sup> The two *pol* gene alignments cover almost whole PR and part of RT (nucleotides 2316–3316 relative to HXB2) and whole INT (nucleotides 4230–5096 relative to HXB2), while the *env* gene alignment covers *gp120-C2V3* region (nucleotides 6905–7288 relative to HXB2). Phylogenetic analyses were performed by the Neighbor-Joining (NJ) algorithm using the Tamura-Nei nucleotide substitution model,<sup>27</sup> in 1000 bootstrap replicates, as implemented in MEGA v4 program.<sup>28</sup> Analysis of recombination was performed by bootscanning analysis (sliding window of 250 bp, incremental steps of 10 bases, and the Kimura two-parameter model<sup>29</sup>) using Simplot 3.5.1 software.<sup>30</sup> Bootstrap support was calculated based on 100 re-samplings.

### 3.4. Resistance analysis

Drug-resistance mutations (DRMs) profile and antiretroviral (ARV) susceptibility were inferred by submitting the PR/RT sequences to the Calibrated Population Resistance Tool (CPR) (version 5.0 beta; available at <http://cpr.stanford.edu/cpr/servlet/CPR>) and the INT sequences to the Stanford HIV Resistance Database (<http://hivdb.stanford.edu/>). The CPR program provides a standard

approach to estimating the prevalence of transmitted HIV drug resistance using population-sampled sequence data and disregard natural occurring polymorphisms and mutations occurring in a frequency higher than 0.5% in untreated subjects.<sup>31,32</sup>

### 3.5. Statistical analysis

The Chi-squared test was employed to evaluate possible association of HIV-1 subtype with demographic, exposure category and clinical variables. A *p* value <0.05 was considered significant.

## 4. Results

Out of 82 HIV-1 isolates analyzed at the PR/RT, INT and *env* regions, 54 (65.8%) were classified as subtype C, 15 (18.3%) as B/C inter-subtype recombinants, 11 (13.5%) as subtype B, one (1.2%) as subtype F1 and one (1.2%) as B/C/F mosaic form (Fig. 1). Three (3.6%) BC recombinant sequences displayed a CRF31\_BC-like recombination profile at PR/RT and clustered among CRF31\_BC reference sequences in the *pol* (PR/RT) NJ tree (Fig. 2). One of these sequences (BR10SC93) displayed one additional B/C recombination point at INT and was classified as a unique recombinant form (URF), as well as the remaining inter-subtype recombinant sequences (Fig. 3).

Table 1 summarizes the demographic characteristics, exposure category, clinical and genetic HIV subtype data of the 82 patients analyzed. Forty (48.8%) patients were males and 42 (51.2%) were females, with a mean age of 37.8 years. The major exposure category was heterosexual (70.6%), followed by men who have sex with men (MSM) (22%) and injection drug users (5%). The herein presented data shows a highly significant association (*p* < 0.001) between exposure category and HIV-1 subtype. Eighty percent of

**Table 2**

Primary drug resistance mutations associated to protease, reverse transcriptase and integrase genes among treatment naïve patients.

Sample	Pol subtype	PI resistance mutations	NRTI resistance mutations	NNRTI resistance mutations	INTI resistance mutations	Resistance profile		
						Low	Intermediate	High
BR10SC06	C	N88D	K219Q	K101E, V106M, G190A, M230L	S147R + E138A	RAL, ELV	NFV	DLV, EFV, NVP
BR10SC07	C					AZT		
BR10SC29	B	M46I	V75M		DDI	NFV		
BR10SC42	C						d4T	
BR10SC46	B							
BR10SC47	C							
BR10SC50	C	M41L, T215C	ABC, DDI, TDF	AZT, d4T				
BR10SC107	C	M41L	G190A	AZT, d4T, ETR	EFV	NVP		
BR10SC131	C		K103N			DLV, EFV, NVP		

PI, protease inhibitors; NRTI, nucleoside reverse transcriptase inhibitors; NNRTI, non-nucleoside reverse transcriptase inhibitors; INTI, integrase inhibitors; ABC, abacavir; AZT, zidovudine; DDI, didanosine; DLV, delavirdine; d4T, stavudine; EFV, efavirenz; ETR, etravirine; NFV, nelfinavir; NVP, nevirapine; TDF, tenofovir; RAL, raltegravir; ELV, elvitegravir.

BC recombinants (URFs\_BC + CRF31\_BC-like) and 76% of subtype C sequences were amplified from individuals of the heterosexual exposure category, whereas 73% of subtype B sequences were in MSM exposure category (Table 1). Subtype B sequences represent 44% of viruses circulating in the MSM population, but only 5% of viruses circulating among heterosexual or female populations. No statistically significant differences at CD4 T cell count, RNA viral load or mean age were found among HIV-1 subtypes.

Among 82 HIV-1 *pol* sequences obtained from drug naïve patients, the presence of at least one DRM was observed in 9 (11.0%) patients (Table 2). DRMs to protease inhibitors (PI) were observed in two (2.4%) patients. Polymorphisms and mutations at protease secondary codons were seen in 17 (20.7%) sequences, being L101/V and T74S the most frequently found (data not shown). Mutations related with resistance to nucleoside reverse transcriptase inhibitors (NRTI) were observed in four (5%) sequences, and M41L was the most common. DRMs to non-nucleoside reverse transcriptase inhibitors (NNRTI) were observed in three (3.6%) of the sequences analyzed, and G190A was the most common. Resistance to integrase inhibitors (INTI) was observed in only one (1.2%) patient who displayed the resistance mutations S147R and E138A, which confer a low-level of resistance to raltegravir and elvitegravir. Minor drug resistance mutations at INT were observed in 11 (13.5%) sequences, being V151I the most frequently found, especially among subtype B samples (data not shown).

One patient showed DRMs to more than one class of antiretroviral drugs. Among the 11 HIV-1 subtype B *pol* sequences, two (18%) displayed DRM, whereas among the 56 HIV-1 subtype C *pol* sequences, eight (14.3%) showed at least one DRM. Nevertheless, that difference was not significant ( $p > 0.05$ ) and no correlation between resistance mutations and HIV-1 subtypes at *pol* was found.

## 5. Discussion

The present work describes a high prevalence (65.8%) of HIV-1 subtype C in Florianópolis metropolitan area, consistent with two recent studies which described a subtype C prevalence of 64% and 79% in other cities of Santa Catarina State.<sup>13,16</sup> This confirms the higher prevalence of subtype C in Santa Catarina when compared with other Southern Brazilian states: 20–30% in Paraná<sup>15,17</sup> and 27–42% in Rio Grande do Sul.<sup>12,13</sup> The molecular epidemiology scenario of Florianópolis also shows a high frequency (18.3%) of BC recombinants. Two out of 15 BC recombinant sequences identified showed a CRF31\_BC-like recombinant pattern at *pol* (PR/RT/INT) and *env* genes, reinforcing that HIV-1 CRF31\_BC circulates at low frequency in Santa Catarina, consistent with previous studies.<sup>13,16</sup> One URF\_BC identified herein displayed a CRF31\_BC-like pattern at PR/RT, but one additional B/C breakpoint at INT, while others URFs\_BC displayed one coincident recombination breakpoint with CRF31\_BC at PR/RT, but larger subtype B fragments. These variants may be second generation recombinants originated from recombination between CRF31\_BC and subtype B strains, as was recently described in Rio Grande do Sul.<sup>25</sup> Although Florianópolis receives a very important number of tourists from Argentina and Uruguay, we found no evidence of circulation of BF recombinant forms commonly observed in those countries (such as CRF12\_BF and CRF38\_BF)<sup>33</sup>, pointing to a minor impact of international tourism on the HIV molecular epidemiology scenario of Florianópolis.

A significant association was observed between HIV-1 subtype and exposure category, suggesting that two independent epidemics are occurring in Santa Catarina. It seems that HIV-1 subtype C broke up the barriers that separate the exposure categories and circulates at high prevalence within both heterosexual (70.7%) and MSM (50%) groups. By contrast, HIV-1 subtype B circulates at high prevalence within MSM group (44.4%), but not within heterosexual

group (5.2%). On the other hand, the high prevalence of URFs\_BC reflects frequent episodes of coinfections and/or superinfections in the studied population. The connection between subtypes B and C epidemics seems to be more frequent in heterosexual population, but further investigations with a higher number of MSM patients are necessary to confirm this observation.

Comparison of the HIV-1 *env* gene results from the current work with those obtained in a previous study performed in the same area<sup>14</sup> revealed a significant increase in the proportion of subtype C at heterosexual population from 56% in 2004 to 83% in 2008/2009; while the proportion of *env* subtype B decreases from 38% to only 15% in the same period. This result suggests a faster spread of subtype C than subtype B among heterosexual population over last years in Santa Catarina. These observations may reveal a different efficacy of homosexual/heterosexual transmission of subtypes B and C, or may reflect different entries and dissemination networks of these subtypes in Santa Catarina. The high proportion of subtype B among females in Rio Grande do Sul and Paraná (38–42%),<sup>12,17</sup> and the recent expansion of subtype C among MSM in São Paulo (Brazil),<sup>34</sup> strongly favor the second hypothesis.

Our findings indicate a prevalence of PR/RT primary DRMs (10%) higher than the national average (6–8%),<sup>2,18,19</sup> although lower than those described in other Brazilian regions (18–37%).<sup>20–22</sup> Notwithstanding, the prevalence of primary resistance presented herein may be underestimated due mostly selected patients were chronic cases (>12 months since HIV infection), and HIV tends to return to wild type over time without ARV selective pressure as DRMs reduce the viral fitness.<sup>35</sup> We also observed one patient with low-level resistance to INTI. Such resistance mutations, however, should not be considered a case of transmitted resistance since the diagnostic date of that patient was 2002, several years before the introduction of Raltegravir in Brazil (2009). Those mutations probably reflect rare substitutions in the integrase gene as described elsewhere.<sup>36–38</sup> These results encourage the use of this new drug in the treatment of AIDS patients in Southern Brazil.

In conclusion, our study indicates an increase of HIV-1 subtype C prevalence in the metropolitan area of Florianópolis with a significantly association between virus subtypes and exposure categories. The Southern region shows the highest incidence of HIV-1 infections in the country, which combined with the increasing contribution of HIV-1 infections among heterosexuals, might explain the expansion of subtype C in the Brazilian epidemic. This combination makes southern Brazil an interesting site for development of health innovations and vaccine trials.<sup>39</sup> The presented work also describes a prevalence of DRMs above the national average, pointing to the importance of sustained surveillance studies in this region.

## Conflict of interest

Authors declare no conflict of interest.

## Acknowledgments

The authors are grateful to the Medical Staff for support during recruitment, to Patrícia H. Stoco and Aline D. Schlindwein for helping with sequencing, and to Fundação de Apoio à Pesquisa Científica e Tecnológica do Estado de Santa Catarina (FAPESC) for financial support.

## References

1. Boletim epidemiológico, AIDS & DST. Brasil: Ministério da Saúde; 2010.
2. Couto-Fernandez JC, Morgado MG, Bongertz V, Tanuri A, Andrade T, Brites C, et al. HIV-1 subtyping in Salvador, Bahia, Brazil: a city with African sociodemographic characteristics. *J Acquir Immune Defic Syndr* 1999;22(3):288–93.



3. Bongertz V, Bou-Habib DC, Brigido LF, Caseiro M, Chequer PJ, Couto-Fernandez JC, et al. HIV-1 diversity in Brazil: genetic, biologic, and immunologic characterization of HIV-1 strains in three potential HIV vaccine evaluation sites. Brazilian Network for HIV isolation and characterization. *J Acquir Immune Defic Syndr* 2000;**23**(2):184–93.
4. Brindeiro RM, Diaz RS, Sabino EC, Morgado MG, Pires IL, Brigido L, et al. Brazilian Network for HIV Drug Resistance Surveillance (HIV-BResNet): a survey of chronically infected individuals. *AIDS* 2003;**17**(7):1063–9.
5. Gadelha SR, Shindo N, Cruz JN, Morgado MG, Galvao-Castro B. Molecular epidemiology of human immunodeficiency virus-1 in the state of Ceara, Northeast, Brazil. *Mem Inst Oswaldo Cruz* 2003;**98**(4):461–3.
6. Cabral VP, Cunha CB, Magalhaes EF, Pinto-Neto LF, Couto-Fernandez JC, Dietze R, et al. Human immunodeficiency virus type-1 subtypes of infected patients in Espirito Santo, Brazil. *Mem Inst Oswaldo Cruz* 2006;**101**(8):881–5.
7. Veras NM, Veras VS, Ramalho ED, Kyaw C, Silva RR, Brigido MM, et al. HIV type 1 genetic variability in central Brazil. *AIDS Res Hum Retroviruses* 2007;**23**(12):1481–90.
8. Cardoso LP, Queiroz BB, Stefani MM. HIV-1 pol phylogenetic diversity and antiretroviral resistance mutations in treatment naive patients from Central West Brazil. *J Clin Virol* 2009;**46**(2):134–9.
9. Machado LF, Ishak MO, Vallinoto AC, Lemos JA, Azevedo VN, Moreira MR, et al. Molecular epidemiology of HIV type 1 in northern Brazil: identification of subtypes C and D and the introduction of CRF02\_AG in the Amazon region of Brazil. *AIDS Res Hum Retroviruses* 2009;**25**(10):961–6.
10. Soares EA, Santos RP, Pellegrini JA, Sprinz E, Tanuri A, Soares MA. Epidemiologic and molecular characterization of human immunodeficiency virus type 1 in southern Brazil. *J Acquir Immune Defic Syndr* 2003;**34**(5):520–6.
11. Rodrigues R, Scherer LC, Oliveira CM, Franco HM, Sperhake RD, Ferreira JL, et al. Low prevalence of primary antiretroviral resistance mutations and predominance of HIV-1 clade C at polymerase gene in newly diagnosed individuals from south Brazil. *Virus Res* 2006;**116**(1–2):201–7.
12. Santos AF, Schrago CG, Martinez AM, Mendoza-Sassi R, Silveira J, Sousa TM, et al. Epidemiologic and evolutionary trends of HIV-1 CRF31\_BC-related strains in southern Brazil. *J Acquir Immune Defic Syndr* 2007;**45**(3):328–33.
13. Brigido LF, Nunes CC, Oliveira CM, Knoll RK, Ferreira JL, Freitas CA, et al. HIV type 1 subtype C and CB Pol recombinants prevail at the cities with the highest AIDS prevalence rate in Brazil. *AIDS Res Hum Retroviruses* 2007;**23**(12):1579–86.
14. Locatelli D, Stoco PH, de Queiroz AT, Alcantara LC, Ferreira LG, Zanetti CR, et al. Molecular epidemiology of HIV-1 in Santa Catarina State confirms increases of subtype C in Southern Brazil. *J Med Virol* 2007;**79**(10):1455–63.
15. Ferreira JL, Thomaz M, Rodrigues R, Harrad D, Oliveira CM, Oliveira CA, et al. Molecular characterisation of newly identified HIV-1 infections in Curitiba, Brazil: preponderance of clade C among males with recent infections. *Mem Inst Oswaldo Cruz* 2008;**103**(8):800–8.
16. Rodrigues R, Manenti S, Romao PR, de Paula Ferreira JL, Batista JP, Siqueira AF, et al. Young pregnant women living with HIV/AIDS in Criciuma, Southern Brazil, are infected almost exclusively with HIV type 1 clade C. *AIDS Res Hum Retroviruses* 2010;**26**(3):351–7.
17. Toledo PV, de Carvalho DS, de Rossi SG, Brindeiro R, de Queiroz-Telles F. Genetic diversity of human immunodeficiency virus-1 isolates in Parana, Brazil. *Braz J Infect Dis* 2010;**14**(3):230–6.
18. Inocencio LA, Pereira AA, Sucupira MC, Fernandez JC, Jorge CP, Souza DF, et al. Brazilian Network for HIV Drug Resistance Surveillance: a survey of individuals recently diagnosed with HIV. *J Int AIDS Soc* 2009;**12**(1):20.
19. Sprinz E, Netto EM, Patelli M, Lima JS, Furtado JJ, da Eira M, et al. Primary antiretroviral drug resistance among HIV type 1-infected individuals in Brazil. *AIDS Res Hum Retroviruses* 2009;**25**(9):861–7.
20. Sucupira MC, Caseiro MM, Alves K, Tescarollo G, Janini LM, Sabino EC, et al. High levels of primary antiretroviral resistance genotypic mutations and B/F recombinants in Santos, Brazil. *AIDS Patient Care STDS* 2007;**21**(2):116–28.
21. de Sa-Filho DJ, Ambar RF, Duarte NB, Matias RB, Candido V, Gagliani LH, et al. HIV type 1 diversity from newly diagnosed patients in Santos metropolitan area/Brazil. *AIDS Res Hum Retroviruses* 2009;**25**(9):925–9.
22. Pedroso C, Queiroz AT, Alcantara LC, Drexler JF, Diaz RS, Weyll N, et al. High prevalence of primary antiretroviral resistance among HIV-1-infected adults and children in Bahia, a northeast state of Brazil. *J Acquir Immune Defic Syndr* 2007;**45**(2):251–3.
23. Delwart EL, Herring B, Rodrigo AG, Mullins JL. Genetic subtyping of human immunodeficiency virus using a heteroduplex mobility assay. *PCR Methods Appl* 1995;**4**(5):S202–16.
24. Bello G, Passaes CP, Guimaraes ML, Lorete RS, Matos Almeida SE, Medeiros RM, et al. Origin and evolutionary history of HIV-1 subtype C in Brazil. *AIDS* 2008;**22**(15):1993–2000.
25. Passaes CP, Bello G, Lorete RS, Matos Almeida SE, Junqueira DM, Veloso VG, et al. Genetic characterization of HIV-1 BC recombinants and evolutionary history of the CRF31\_BC in Southern Brazil. *Infect Genet Evol* 2009;**9**(4):474–82.
26. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 1994;**22**(22):4673–80.
27. Tamura K, Nei M. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol* 1993;**10**(3):512–26.
28. Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol* 2007;**24**(8):1596–9.
29. Kimura M. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 1980;**16**(2):111–20.
30. Lole KS, Bollinger RC, Paranjape RS, Gadkari D, Kulkarni SS, Novak NG, et al. Full-length human immunodeficiency virus type 1 genomes from subtype C-infected seroconverters in India, with evidence of intersubtype recombination. *J Virol* 1999;**73**(1):152–60.
31. Bennett DE, Camacho RJ, Otelea D, Kuritzkes DR, Fleury H, Kiuchi M, et al. Drug resistance mutations for surveillance of transmitted HIV-1 drug-resistance: 2009 update. *PLoS One* 2009;**4**(3):e4724.
32. Shafer RW, Rhee SY, Pillay D, Miller V, Sandstrom P, Schapiro JM, et al. HIV-1 protease and reverse transcriptase mutations for drug resistance surveillance. *AIDS* 2007;**21**(2):215–23.
33. Bello G, Aulicino PC, Ruchansky D, Guimarães ML, Lopez-Galindez C, Casado C, et al. Phylodynamics of HIV-1 circulating recombinant forms 12.BF and 38.BF in Argentina and Uruguay. *Retrovirology* 2010;**22**(3):7–22.
34. Brigido LF, Ferreira JL, Almeida VC, Rocha SQ, Ragazzo TG, Estevam DL, et al. Type 1 C expansion into the State of São Paulo, Brazil. *AIDS Res Hum Retroviruses* 2010;**26**(00):1–6.
35. Payne BA, Nsutebu EF, Hunter ER, Olarinde O, Collini P, Dunbar JA, et al. Low prevalence of transmitted antiretroviral drug resistance in a large UK HIV-1 cohort. *J Antimicrob Chemother* 2008;**62**(3):464–8.
36. Lataillade M, Chiarella J, Kozal MJ. Natural polymorphism of the HIV-1 integrase gene and mutations associated with integrase inhibitor resistance. *Antivir Ther* 2007;**12**(4):563–70.
37. Rhee SY, Liu TF, Kiuchi M, Zioni R, Gifford RJ, Holmes SP, et al. Natural variation of HIV-1 group M integrase: implications for a new class of antiretroviral inhibitors. *Retrovirology* 2008;**5**(8):74.
38. Ceccherini-Silberstein F, Malet I, Fabeni L, Dimonte S, Svicher V, D'Arrigo R, et al. Specific HIV-1 integrase polymorphisms change their prevalence in untreated versus antiretroviral-treated HIV-1-infected patients, all naive to integrase inhibitors. *J Antimicrob Chemother* 2010;**65**(11):2305–18.
39. Ersching J, Pinto AR. HIV-1 vaccine clinical trials: the Brazilian experience. *Rev Med Virol* 2009;**19**(5):301–11.